

Product Description

Conquer™ Ready-To-Use Disinfectant Cleaner is a cleaner, disinfectant, deodorizer, bactericide, virucide, tuberculocide, fungicide and mildestat. This product is effective against a broad spectrum of pathogenic microorganisms including bacteria, antibiotic resistant bacteria, viruses, fungi, mold, and mildew.

Conquer™ Ready-To-Use Disinfectant Cleaner is designed specifically as a general non-acid cleaner and disinfectant for use on hard, non-porous surfaces in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food.

Packaging

1 quart bottle

12 bottles per case

12 trigger sprayer applicators per case

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces.

DISINFECTION, DEODORIZING AND CLEANING – Remove visible soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 1 minute). No scrubbing is necessary. Wipe dry with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

ONE-STEP* CLEANING AND DISINFECTION OF TOILET BOWLS AND URINALS: Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of this product evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 1 minute) and flush. If water cannot be removed from toilet, use product on the external surface and above waterline only.

BACTERICIDAL ACTIVITY: Using AOAC test methods (under Good Laboratory Practices, [GLPs], in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, this product kills the following organisms on hard non-porous inanimate surfaces: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella (choleraesuis) enterica*, *Escherichia coli*, *Escherichia coli O157:H7*, *Streptococcus pyogenes (Necrotizing Fasciitis-Group A)* (V.A. Medical Center Isolate 04001), *Listeria monocytogenes*, *Yersinia enterocolitica*, *Enterococcus faecium*, *Corynebacterium ammoniagenes*, *Salmonella (typhi) enterica*.

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Using AOAC test methods (under Good Laboratory Practices, [GLPs], in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, this product kills the following organisms on hard non-porous inanimate surfaces: Methicillin resistant *Staphylococcus aureus* (MRSA), Vancomycin resistant *Enterococcus faecalis* (VRE), Vancomycin intermediate resistant *Staphylococcus aureus* (VISA) (CDC Isolate 99287), Methicillin resistant *Staphylococcus epidermidis* (MRSE), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) (NRS 123) Genotype USA400, Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) (NRS 384) Genotype USA300.

TUBERCULOCIDAL ACTIVITY - This product exhibits disinfectant efficacy against *Mycobacterium tuberculosis* BCG (*Mycobacterium bovis*) at 20 degrees Centigrade when the treated surface is allowed to remain wet for 5 minutes.

FUNGICIDAL ACTIVITY – This product is fungicidal against the pathogenic fungus, *Trichophyton interdigitale*, formerly *Trichophyton mentagrophytes*, a cause of ringworm of the foot, when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms or other areas commonly contacted by bare feet.

MILDEWSTAT – To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

†VIRUCIDAL ACTIVITY – This product kills on hard, non-porous inanimate surfaces when allowed to remain wet for a 10 minute contact time against: Canine Parvovirus, Hepatitis A Virus and Poliovirus Type 1. This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: Human Coronavirus, SARS associated Coronavirus, Avian Influenza A strain H3N2 and Avian Influenza A strain H9N2 for 2 minutes; Paramyxovirus (Mumps), Rhinovirus type 39 and Rotavirus for 3 minutes; HIV-1 (associated with AIDS), SARS Related Coronavirus 2 [SARS-CoV-2] [USA-WA1/2020] [BEI NR-52281] for 1 minute; Hepatitis B Virus (HBV), Hepatitis C Virus (HCV) and Bovine Viral Diarrhea Virus (BVDV) (NVSL) when the treated surface is allowed to remain wet for 1 minute and Norovirus (Norwalk virus), Feline Calicivirus and Rabies Virus for 30 seconds.

KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS

in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV and HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface to remain wet for HBV and HCV 1 minute, for HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

Storage And Disposal

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. **PESTICIDE STORAGE** - Store in a dry place no lower in= temperature than 50°F or higher than 120°F. **PESTICIDE DISPOSAL** - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL** - Nonrefillable container. Do not reuse or refill container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.

VIRUCIDAL - 30 SECOND KILL

Norovirus (Norwalk virus)
Feline Calicivirus (ATCC VR-782)
Rabies Virus

VIRUCIDAL - 1 MINUTE KILL

Pathogen Contact Time Hepatitis B Virus (HBV)
Hepatitis C Virus (HCV)
Bovine Viral Diarrhea Virus (BVDV)
HIV-1 (associated with AIDS)

**SARS-Related Coronavirus 2 (SARS-CoV-2)
USA-WA1/2020 BEI NR-52281**

BACTERICIDAL - 3 MINUTE KILL

Pseudomonas aeruginosa (ATCC 15442)
Staphylococcus aureus (ATCC 6538)
Salmonella (choleraesuis) enterica (ATCC 10708)
Escherichia coli (ATCC 11229)
Escherichia coli O157:H7 (ATCC 43895)
Streptococcus pyogenes (Necrotizing Fasciitis Group A) (V. A. Medical Center Isolate 04001)
Listeria monocytogenes (ATCC 35152)
Yersinia enterocolitica (ATCC 23715)
Enterococcus faecium (ATCC 6569)
Corynebacterium ammoniagenes (ATCC 6871)
Salmonella (typhi) enterica (ATCC 6539)
Methicillin-resistant Staphylococcus aureus (MRSA) (ATCC 33593)
Vancomycin-resistant Enterococcus faecalis (VRE) (ATCC 51575)
Vancomycin intermediate-resistant Staphylococcus aureus (VISA) (CDC Isolate 99287)
Methicillin-resistant Staphylococcus epidermidis (MRSE) (ATCC 51625)
Community Associated Methicillin-resistant Staphylococcus aureus (CA-MRSA) (NRS 123) Genotype USA 400
Community Associated Methicillin-resistant Staphylococcus aureus (CA-MRSA) (NRS 384) Genotype USA 300

VIRUCIDAL - 2 MINUTE KILL

Human Coronavirus (ATCC VR-740)
SARS associated Coronavirus
Avian Influenza A Strain H3N2 (ATCC VR-2072)
Avian Influenza A Strain H9N2 \

VIRUCIDAL - 3 MINUTE KILL

Paramyxovirus (Mumps) (ATCC VR-1438)
Rhinovirus type 39 (ATCC VR-340)
Rotavirus (ATCC VR-2018)

VIRUCIDAL - 10 MINUTE KILL

Canine parvovirus (ATCC VR-2017)
Hepatitis A Virus (HAV) (University of Ottawa)
Poliovirus Type 1 (ATCC VR-1000)

TUBERCULOCIDAL - 5 MINUTE KILL

Mycobacterium bovis

FUNGICIDAL - 10 MINUTE KILL

Trichophyton interdigitale (ATCC 9533)
--



Eliminates 99.9%* of the virus that causes COVID-19** on hard, non-porous surfaces in 1 minute!

*when used according to disinfection directions
**SARS-Related Coronavirus 2